



Product presentation



Product characteristics

- **Active ingredient:** metoclopramide (as hydrochloride)
- **Pharmaceutical form:** nasal spray solution
- **Route of administration:** nasal use
- **Strength:** 10 mg/ dose and 20 mg/dose
- **Packaging:** BT x 1 bottle x 6ml + dosing pump



Therapeutic indications

Adult population

- Prevention of post-operative nausea and vomiting (PONV)
- Prevention of delayed chemotherapy induced nausea and vomiting (CINV)
- Prevention of radiotherapy induced nausea and vomiting (RINV)
- Symptomatic treatment of nausea and vomiting, including acute migraine nausea and vomiting. Metoclopramide can be used in combination with oral analgesics for the improvement of the absorption of the analgesics in acute migraine



Therapeutic indications

Paediatric population

The product is indicated for children aged 1-18 years for:

- Prevention of delayed chemotherapy induced nausea and vomiting (CINV) as a second line option



Dosage

10 mg/dose

- 1 spray of 10 mg in each nostril, 2-3 times per day (dose of 20 – 60 mg).
Administration is recommended before meal.

20 mg/dose

- 1 spray of 20 mg in one nostril, 2-3 times per day (dose of 40 – 60 mg).
Administration is recommended before meal.