



verisfield®

**Product presentation**



**Xylocream®**

verisfield®

## **Product characteristics**

- **Active ingredients:** Lidocaine and Prilocaine
- **Pharmaceutical form:** Cream
- **Route of administration:** Topical use
- **Strength:** (2,5% + 2,5%) w/w
- **Packaging:** 1 tube x 30 g



**Xylocream®**

verisfield®

## **Indications - usage**

- Topical anaesthesia of the intact skin connected to superficial surgical interventions and dermal procedures, e.g.:
  - needle insertion & surgical treatment of localized lesions
  - curettage of molluscum contagiosum
  - split-skin grafting
  - punch or excision skin biopsy
  - laser treatment of facial port wine stains, hair removal, waxing
  - mesotherapy, fillers



**Xylocream®**

verisfield®

## **Indications - usage**

- Topical anaesthesia of the genital mucosa/genital area connected to:
  - removal of condylomata acuminata
  - lesions of vulval mucosa membranes in females
  - prior to superficial surgical procedures or prior to infiltration anaesthesia of mucosa
  - laser hair removal/waxing



**Xylocream®**

verisfield®

## **Indications - usage**

- Surface anaesthesia of leg ulcers to facilitate mechanical cleansing/debridement
- In newborn, infants and children under the age of 18 years for the local anaesthesia on intact skin prior to minor dermatological procedures (e.g. curettage of molluscum contagiosum) or prior to needle insertion



**Xylocream<sup>®</sup>**

verisfield<sup>®</sup>

## **Mechanism of action/pharmacodynamics - pharmacokinetics**

- Lidocaine and prilocaine are both amide-type local anaesthetic agents that inhibit nerve conduction by interfering with the entry of sodium ions through the nerve cell membrane
- Xylocream<sup>®</sup> provides dermal analgesia. The depth of analgesia depends upon the application time and the dose. It causes transient local peripheral vasoconstriction or vasodilation at the treated area
- In genital mucosa, absorption is more rapid compared to skin application



**Xylocream®**

verisfield®

## **Xylocream® characteristics**

- Easy to use topical anaesthetic. Depending on the dose and the duration of application it can provide up to 2 h of anesthesia after removal of the cream
- Safe to use in neonates, infants, children and adults
- Useful for almost all superficial dermal procedures (medical aesthetic), genital mucosa and the genital skin area procedures, leg ulcers cleansing